

# A Webbing Barklouse, A Psocid, *Archipsocus nomas* Gurney (Insecta: Psocoptera: Archipsocidae)<sup>1</sup>

Donald W. Hall<sup>2</sup>

## Introduction

*Archipsocus nomas* Gurney is a communal web-spinning barklouse. During some years they make extensive silken webs that often cover the trunks and branches of trees in the southeastern US. The webs are believed to protect the barklice from predators. The webs are unsightly, but neither the barklice nor the webs cause any harm to the trees. These insects are sometimes referred to as “tree cattle.”

## Distribution

Webbing barklice are found along the Gulf coast from Texas to Florida and along the Atlantic coast north to South Carolina.

## Description

The eggs of *A. nomas* are oblong, wider at one end, and gray or white in color. Nymphs and adults are soft-bodied insects with chewing mouthparts and long filamentous antennae. They lack cerci. First instar nymphs have eight antennal segments, but the full complement of 13 segments is gained at the first molt. The number of nymphal instars has not been determined for *A. nomas*. In *Archipsocus floridanus*, a closely related species, females have six, and males may have from four to six—but most commonly five. Adults are darker in color than nymphs. When winged, the wings are membranous and held roof-like over the

body. Winged females may be either long or short winged. Winged males are short winged.



Figure 1. Young *Archipsocus* sp. nymphs.  
Credits: Lyle Buss, UF/IFAS



Figure 2. Late instar *Archipsocus* sp. nymphs.  
Credits: Lyle Buss, UF/IFAS

1. This document is EENY275, one of a series of the Entomology and Nematology Department, UF/IFAS Extension. Original publication date October 2002. Revised November 2005 and March 2016. Reviewed April 2019. Visit the EDIS website at <https://edis.ifas.ufl.edu> for the currently supported version of this publication. This document is also available on the Featured Creatures website at <http://entomology.ifas.ufl.edu/creatures>.

2. Donald W. Hall, Entomology and Nematology Department; UF/IFAS Extension, Gainesville, FL 32611.

The Institute of Food and Agricultural Sciences (IFAS) is an Equal Opportunity Institution authorized to provide research, educational information and other services only to individuals and institutions that function with non-discrimination with respect to race, creed, color, religion, age, disability, sex, sexual orientation, marital status, national origin, political opinions or affiliations. For more information on obtaining other UF/IFAS Extension publications, contact your county's UF/IFAS Extension office.

U.S. Department of Agriculture, UF/IFAS Extension Service, University of Florida, IFAS, Florida A & M University Cooperative Extension Program, and Boards of County Commissioners Cooperating. Nick T. Place, dean for UF/IFAS Extension.



Figure 3. *Archipsocus* sp. winged females.  
Credits: Lyle Buss, UF/IFAS

## Life Cycle and Biology

Colonies of *A. nomas* are rare during the winter (December–February in central Florida) and are concentrated in live oak (*Quercus virginiana* Mill.) hammocks and stands of cabbage palms (*Sabal palmetto* [Walter]) where they probably are best sheltered from killing frosts. During this time, colonies are composed primarily of adults with a few late instar nymphs. In the spring, eggs are laid singly or in groups and covered with debris or sometimes by feces. From March to June, colonies become more frequent, and from July to October, colonies rapidly increase in both number and size. The long-winged females are usually observed only during this time of maximum colony proliferation. Nymphs and adults probably feed primarily on lichens. By early December, populations have been greatly reduced by frost, and the webs begin to disintegrate due to weathering. Because the barklice are not harmful, control measures are not recommended.



Figure 4. *Archipsocus nomas* Gurney silken web on *Ilex* sp.  
Credits: D. W. Hall, UF/IFAS



Figure 5. Tree trunk covered in webbing produced by *Archipsocus nomas* Gurney.  
Credits: Douglas L. Caldwell, UF/IFAS

## Selected References

- Borror DJ, Triplehorn CA, Johnson NF. 1989. *An Introduction to the Study of Insects*. Philadelphia, PA: Saunders College Publishing.
- Mockford EL. 1953. "Three new species of *Archipsocus* from Florida (Psocoptera: Archipsocidae)." *Florida Entomologist* 26: 113–124.
- Mockford EL. 1957. Life history studies on some Florida insects of the genus *Archipsocus* (Psocoptera). Gainesville, FL: Bulletin of the Florida State Museum (Biological Sciences) University of Florida. 1: 253–274.
- Mockford EL. 1987. Order Psocoptera. In: Stehr FW, ed., *Immature Insects*. Dubuque, IA: (Vol. 1) Kendall/Hunt Publishing Company.